

Graduated compression stockings in the prevention of venous thromboembolism

O. Agu, G. Hamilton and D. Baker

University Department of Surgery, Royal Free Hospital, Pond Street, Hampstead, London NW3 2QP, UK

Correspondence to: Mr O. Agu

Background: Surveys still show a wide variation in routine use of deep vein thrombosis (DVT) prophylaxis despite its established place in current patient management. This article reviews the mechanism of action, efficacy and complications of stockings in preventing DVT.

Methods: Relevant publications indexed in Medline (1966–1998) and the Cochrane database were identified. Appropriate articles identified from the reference lists of the above searches were also selected and reviewed.

Results and conclusion: Graduated compression stockings reduce the overall cross-sectional area of the limb, increase the linear velocity of venous flow, reduce venous wall distension and improve valvular function. Fifteen randomized controlled trials of graduated compression stockings alone were reviewed. Stockings reduced the relative risk of DVT by 64 per cent in general surgical patients and 57 per cent following total hip replacement. The effect of stockings was enhanced by combination with pharmacological agents such as heparin; the combination is recommended in patients at moderate or high risk of DVT. Knee-length stockings are as effective and should replace above-knee stockings. Complications are rare and avoidable.

Paper accepted 18 April 1999

British Journal of Surgery 1999, 86, 992–1004

Prevention of deep vein thrombosis (DVT) in hospitalized patients has received considerable attention because of the risk of potentially fatal pulmonary thromboembolism and the long-term morbidity associated with chronic venous valvular insufficiency. The result has been a continuing drop in the incidence and complications of DVT^{1–5}. Without prophylaxis, the risk of DVT ranges from 25 to 30 per cent in general surgical patients to 70 per cent following some orthopaedic procedures^{6–9}. Prophylaxis, often consisting of a combination of pharmacological and mechanical agents, reduces the risk of DVT by 60–80 per cent in general surgical and 50–64 per cent in orthopaedic patients^{10–13}.

Despite the established place of prophylaxis in current patient management, more episodes of venous thromboembolism are reported in clinical practice than would be expected. Surveys show a wide variation in routine use of DVT prophylaxis^{14–17}. This variation may represent a lag between intended and actual implemented prophylaxis. There appears to be a shortfall in awareness of the mechanism of action and efficacy, especially of mechanical

agents. Violation of established protocols, which may involve up to 30 per cent of patients, has also been reported¹⁸.

Graduated compression stockings are the most widely used mechanical antithrombotic agents. They are of proven efficacy in general surgical patients¹¹. However, the case for stockings may not be as convincing in some other surgical specialties, especially orthopaedics, where patients are at much greater risk^{19–21}. The value of stockings for obstetric, long-stay general medical and geriatric patients is also unclear. Evidence supporting current stocking pressure profiles, which have been in use for over 25 years, needs reappraisal especially in patient groups in which stockings are not proving as effective. Evidence comparing the efficacy of thigh- versus knee-length stockings needs clarification. Although stockings are generally regarded as safe, complications occasionally occur.

This review considers the current evidence on the mechanism of action, design, efficacy and complications associated with graduated compression stockings in the prevention of venous thromboembolism.




Fig. 1 Possible mechanism of action of graduated compression stocking in preventing deep vein thrombosis

Methods

Relevant publications indexed in Medline (1966–1998) were identified using the medical subject headings ‘stockings’, ‘clothing’, ‘bandages’, ‘thrombophlebitis’, ‘venous thrombosis’, ‘postoperative complications’ and ‘thrombosis/prevention’. The Cochrane database was also searched and relevant studies identified. Further articles identified from the reference lists of the above searches were also selected and reviewed. Odds ratios of studies reviewed were combined using the Mantel–Haenszel χ^2 procedure. Number needed to treat was calculated based on a 25 per cent risk of DVT in untreated general surgical patients and a 40 per cent risk following total hip replacement.

Mechanism

The mechanism of action of compression stockings in preventing venous thrombosis remains unclear and is probably multifactorial (Fig. 1). Compression stockings achieve their effect by affecting the three classical aetiological factors described by Virchow (1856), namely stasis, endothelial damage and hypercoagulability. A decrease in the luminal diameter of the vein appears to play the key role.

Effect of external compression on venous function

External compression reduces the overall cross-sectional area of the lower limb and increases the linear velocity of blood flow within the veins^{22–27}. An external pressure of 15 mmHg results in a 20 per cent reduction in the venous cross-sectional area and a significant increase in the velocity of flow in both superficial and deep venous systems²⁸. Increased velocity of flow reduces venous stasis and decreases the risk of thrombus formation by reducing venous wall distension, local contact time and the concentration of coagulation reactants. Compression stockings also improve evacuation of incompetent and incompletely emptied valvular cusps²⁹. Stasis within these cusps is associated with the initiation of venous thrombosis³⁰. Stockings have also been shown to enhance venous function by improving coaptation of valvular cusps, again reducing reflux-related stasis and the risk of thrombosis, especially in patients with valvular insufficiency³¹.

Postoperative DVT is associated with intraoperative systemic venodilatation affecting both lower and upper limbs^{32,33}. Passive venodilatation stretches the endothelium beyond the support of the tunica media. The resulting intimal tears have been demonstrated on electron micro-

scopy in canine models. Intimal tears predispose to DVT by exposing thrombogenic subendothelial collagen, in the presence of venous stasis, to activated platelets, clotting factors and other thrombogenic products of tissue injury³⁴. Venodilatation in excess of 20 per cent of resting diameter has been shown to be associated with an increased risk of DVT³². Coleridge Smith *et al.*³⁵ demonstrated a median 48 (interquartile range 26–53) per cent reduction in the diameter of gastrocnemius veins during operation in patients wearing graduated compression stockings. Additionally, in the same study, in contrast to the control group, no venous distension occurred during operation in patients wearing stockings.

Elastic compression may achieve some of its effects by altering the balance of coagulation reactants in patients at risk of thrombosis. Elastic compression in the supine reverse Trendelenburg position has been shown to increase systemic plasma levels of tissue factor pathway inhibitor (TFPI)³⁶. TFPI is a recently described factor Xa-dependent inhibitor of the tissue factor (extrinsic) coagulation pathway. TFPI is endothelium derived, and increases with shear stress in the upright posture, after extensive exercise and following administration of both unfractionated heparin and low molecular weight heparins (LMWHs)^{37–41}. Increased TFPI in the upright tilt may be an endothelial response to venous stasis and luminal distension to prevent local thrombosis. Enhancing this TFPI effect by elastic compression may represent another mechanism of local and systemic venous thromboprophylaxis.

Design of compression stockings

Compression stockings may be designed to apply graduated or uniform compression. Graduated compression refers to the application of varying degrees of constant compression to different segments of the leg, with pressure being greatest at the ankle and gradually decreasing proximally. Uniform compression aims to deliver a similar degree of pressure to the whole leg. Graduated compression has been the favoured option since Sigel *et al.*⁴² demonstrated small but consistent increases in velocity of flow in the common femoral vein compared with that obtained with uniform compression. This conclusion was based on the effect of a uniform compression of 11 mmHg in four subjects in a supine position at 15° foot-down tilt. The hydrostatic gradient introduced by this position (about 14 mmHg) is scarcely representative of the patient lying in hospital and is large enough to affect the validity of their results. While graduated compression is of proven efficacy in preventing DVT, evidence to support its superiority over uniform compression appears inconclusive and needs revalidation. Discrediting uniform compression on the basis of trials of

Tubigrip (Seton Health Group, Oldham, UK) is probably inappropriate, as the latter is unlikely to deliver uniform compression with increasing proximal lower limb circumference. Instead, a reverse compression gradient or tourniquet effect is to be expected.

The design of the majority of stockings in current use is based on a pressure profile of 18, 14, 8, 10, 8 mmHg from ankle to thigh as recommended by Sigel *et al.*⁴² and Lawrence and Kakkar⁴³, and confirmed by clinical trials in comparison to untreated controls^{44,45}. Although 30 mmHg at the ankle had a greater effect on velocity of flow than 18 mmHg, the latter was deemed safer, being less likely to compromise subcutaneous oxygenation^{43,46}. Sparrow *et al.*⁴⁷ showed that the efficiency of stockings correlated closely with changes in calf compression if ankle compression was held constant; higher calf compression was more effective. Varying ankle compression and ankle–calf gradient were not as important. A profile of 16.8, 14.5 and 6.4 mmHg at the ankle, calf and thigh respectively proved most effective. Although the 18 to 8 mmHg pressure profile is effective, it may not necessarily represent the best profile or be ideal in all clinical situations in which stockings are needed for DVT prophylaxis. Further studies are needed to determine the optimal profile in different clinical situations. These studies would require a large number of subjects, a wide variety of pressure profiles and postures, and should be based on the effect of stockings on both venodilatation and velocity of venous outflow.

Efficacy

Results of initial trials on the value of compression for venous thromboprophylaxis were equivocal. Wilkins *et al.*⁴⁸ reported, in a post-mortem study, one non-fatal pulmonary embolism (PE) in 100 deaths in 2346 patients with stockings compared with six PEs (two fatal) in 89 deaths in 2395 untreated non-randomized controls. Makin⁴⁹ reported a reduced incidence of clinically diagnosed DVT with Tubigrip. Rosengarten *et al.*⁵⁰ and Browse *et al.*⁵¹ were unable to demonstrate any difference between stockings and control using the fibrinogen uptake test. These discrepancies probably resulted from the use of Tubigrip for compression, which exerts non-uniform and potentially reverse gradient compression. In addition, the studies used clinical diagnosis, which is undoubtedly unreliable in predicting the incidence of DVT and PE. Subsequent studies using better stocking design and more reliable diagnostic tools provide a clearer picture.

General surgery

The majority of studies of stockings for the prevention of venous thrombosis have been on general surgical patients.

Table 1 Results of randomized controlled trials of graduated compression stockings in preventing deep vein thrombosis

Reference	Year	Diagnostic test	Incidence of deep vein thrombosis*		Odds ratio†	Risk reduction (%)	No. needed to treat‡
			Stockings	Control			
General/abdominal surgery							
Holford ⁴⁴	1976	FUT	11 of 48 (23)	23 of 47 (49)	0.31 (0.12–0.82)	53	6
Scurr <i>et al.</i> ⁵²	1977	FUT	8 of 70 (11)	26 of 70 (37)	0.22 (0.08–0.57)	69	5
Torngren ⁵³	1980	FUT	4 of 98 (4)	12 of 98 (12)	0.31 (0.08–1.07)	67	6
Allan <i>et al.</i> ⁴⁵	1983	FUT/venography	15 of 97 (15)	37 of 103 (36)	0.33 (0.16–0.68)	57	7
Bergqvist and Lindblad ⁵⁴	1984	FUT	0 of 80 (0)	8 of 80 (10)	0.00 (0.00–0.55)	100	4
Wille-Jorgensen <i>et al.</i> ⁵⁵	1985	FUT/venography	2 of 86 (2)	11 of 90 (12)	0.17 (0.03–0.86)	81	5
Mellbring and Palmer ⁵⁶	1986	FUT	1 of 54 (2)	2 of 54 (4)	0.49 (0.02–7.22)	50	9
			7 of 54 (13)	6 of 54 (11)	1.19 (0.32–0.39)	-17	-50
Scurr <i>et al.</i> ⁵⁷	1987	FUT/venography	1 of 78 (1)	7 of 78 (9)	0.13 (0.01–0.11)	86	5
Wille-Jorgensen <i>et al.</i> ⁵⁸	1991	FUT/venography	2 of 83 (2)	12 of 83 (14)	0.14 (0.02–0.70)	84	5
Total			51 of 748 (7)	144 of 757 (19)	0.31 (0.22–0.44)	64	8
Gynaecology							
Turner <i>et al.</i> ⁵⁹	1984	FUT	0 of 104 (0)	4 of 92 (4)	0.00 (0.00–1.32)	100	7
Neurosurgery							
Turpie <i>et al.</i> ⁶⁰	1989	FUT/venography	7 of 80 (9)	16 of 81 (20)	0.38 (0.14–1.09)	56	7

Values in parentheses are *percentages or †95 per cent confidence intervals. ‡Based on PEER (patient expected event rate) = 0.25. FUT, fibrinogen ¹²⁵I uptake test; venography, contrast venography

Table 1 shows the results of randomized controlled trials comparing the incidence of DVT in general surgical patients. The overall incidence of DVT was 51 (7 per cent) of 748 in the treatment group compared with 144 (19 per cent) of 757 in the control group. The summary odds ratio was 0.31, which translated to a relative risk reduction of 64 (95 per cent confidence interval 53–73) per cent in patients wearing stockings. Treating eight general surgical patients with graduated compression stockings prevented the occurrence of one DVT. Wells *et al.*⁶¹ reported a 68 per cent relative risk reduction of DVT in general surgical patients using stockings. In practice, stockings are often combined with pharmacological agents, especially low-dose heparin or LMWH. This combination leads to improved results (see later).

Direct evidence of the value of graduated compression stockings in preventing fatal PE is inconclusive. Numbers limit most trials, as a large sample size would be required to show a significant reduction because of the low incidence of PE, particularly as the widespread use of heparin has substantially reduced the risk of DVT and PE. The effect of stockings in preventing the long-term post-thrombotic sequelae of DVT is easier to demonstrate. Some 60 per cent of patients treated for DVT will have residual venous valvular insufficiency⁶² with symptoms ranging from leg swelling and discomfort to venous ulcers. Graduated compression stockings have been shown to prevent primary and recurrent DVT, and decrease the risk and severity of the post-thrombotic syndrome^{61,63,64}. A recent randomized clinical trial of the effect of compression stockings in 194

patients following symptomatic proximal vein thrombosis reported mild to moderate post-thrombotic syndrome in 19 patients (20 per cent) wearing stockings compared with 46 controls (47 per cent) ($P < 0.001$). In addition, only 11 patients (11 per cent) wearing stockings developed a severe post-thrombotic syndrome compared with 23 control patients (23 per cent) ($P < 0.001$)⁶⁴.

Orthopaedic surgery

The majority of studies in orthopaedics have included patients undergoing elective total hip replacement. Elderly patients with a fractured hip may constitute a more ideal study population as they are often dehydrated, immobile and have a much greater risk of venous thromboembolism¹³. Orthopaedic surgeons in the UK are more likely to use routine prophylaxis in patients undergoing elective hip replacement than surgery for hip fracture despite a fivefold increased risk of fatal PE in the latter group⁶⁵. The incidence of DVT exceeds 50 per cent in untreated controls (Table 2); however, thromboprophylaxis appears to be less effective in orthopaedic than in other forms of surgery^{10,70–73}. Although the majority of orthopaedic surgeons use pharmacological prophylaxis, the risk of bleeding leading to wound haematoma, infection and implant failure is perceived as unacceptable by a minority⁷⁴. Physical agents are safer and more acceptable in this regard^{75,76}. Examining the results of the randomized trials shows a 57 per cent overall relative risk reduction in incidence of DVT following total hip replacement in patients who used

Table 2 Randomized controlled trials of graduated compression stockings for deep vein thrombosis prophylaxis following total hip replacement

Reference	Year	Diagnostic test	Incidence of deep vein thrombosis*		Odds ratio†	Risk reduction (%)	No. needed to treat‡
			Stockings	Control			
Ishak and Morley ⁶⁶	1981	Venography	7 of 35 (20)	22 of 41 (54)	0.22 (0.07–0.67)	63	4
Fredin <i>et al.</i> ⁶⁷	1989	Venography	13 of 44 (30)	21 of 46 (46)	0.50 (0.19–1.30)	35	7
Kalodiki <i>et al.</i> ⁶⁸	1996	Venography	12 of 38 (32)	13 of 14 (93)	0.04 (0.00–0.32)	66	3
Barnes <i>et al.</i> ⁶⁹	1978	Doppler	0 of 8 (0)	5 of 10 (50)	0.00 (0.00–1.08)	100	2
Total			32 of 125 (26)	61 of 111 (55)	0.23 (0.12–0.42)	57	4

Values in parentheses are *percentages or †95 per cent confidence intervals. ‡Based on PEER (patient expected event rate) = 0.4. Venography, contrast venography; Doppler, Doppler ultrasonography

stockings compared with controls (*Table 2*). Treating only four patients with stockings prevented the occurrence of one DVT following total hip replacement. A recent meta-analysis of methods of prophylaxis following total hip replacement suggested that, while all recommended methods except aspirin decreased the risk, LMWH and stockings had the greatest relative value in preventing venous thromboembolism⁷⁷. Although LMWH proved more effective, it was associated with a risk of bleeding and heparin-induced thrombocytopenia^{41,78–80}. Reporting a placebo-controlled, randomized clinical trial on 78 patients undergoing total hip replacement, Kalodiki *et al.*⁶⁸ demonstrated that the combination of LMWH (enoxaparin 40 mg daily) and stockings was more effective in preventing DVT than LMWH alone, and that stockings alone were better than nothing (13, 28 and 57 per cent incidence of proximal DVT in LMWH and stocking group, LMWH alone and untreated control groups respectively). The combination of LMWH and stockings is recommended in patients undergoing hip replacement.

Venous thromboembolism following total knee replacement is especially resistant to prophylaxis^{79,81,82}. This may be due to intraoperative positioning for optimal access, direct vascular injury from intraoperative limb manipulation and the thigh tourniquet which is inflated to 450–500 mmHg to achieve a bloodless field^{83,84}. Use of a thigh tourniquet impairs venous drainage, damages the endothelium, confines coagulation reactants from surgical trauma below the tourniquet and may increase the risk of venous thrombosis and large emboli^{85–88}. Hui *et al.*¹⁹ reported a significant reduction in proximal and calf venous thrombosis after total knee replacement when patients wore below-knee stockings compared with that in controls (32 *versus* 66 per cent). Above-knee stockings were less effective.

The risk of late-onset venous thrombosis persists for at least 5 weeks after joint replacement^{89,90}. Patients may therefore benefit from continued use of stockings after discharge from hospital. The value of graduated compression

stockings in patients with a fractured hip is still under evaluation.

Gynaecology, obstetrics and urology

The overall risk of DVT ranges between 4 and 45 per cent in gynaecological and 8–80 per cent in urological patients⁹¹, the highest risk being after operations for malignancy⁹². The risk in pregnancy is five times higher than that in non-pregnant age-matched patients. A randomized controlled trial of the value of stockings in 196 patients undergoing major gynaecological surgery showed a significant reduction in venous thrombosis (zero in treated *versus* 4 per cent in controls; $P < 0.05$)⁵⁹. In addition, by preventing venous pooling, stockings may also reduce catecholamine release with beneficial effects for the circulation in pregnancy^{93,94}. Clinical data on the effect of stockings in preventing DVT in pregnancy are, however, insufficient.

There are no prospective randomized controlled studies of the effect of stockings in urological patients. Hansberry *et al.*⁹⁵ compared use of compression stockings with pneumatic compression alone or heparin–dihydroergotamine alone in 74 patients undergoing neoplastic urological procedures using indium-labelled platelet scans. There was no significant difference in the incidence of DVT in the three groups.

Neurology and neurosurgery

Thromboembolism is a major concern in neurological and neurosurgical patients, especially those with spinal cord injury, brain tumour, head injury, stroke and patients undergoing surgery. The risk of DVT in general neurosurgical patients ranges between 19 and 50 per cent⁹⁶. The main advantage of mechanical methods of thromboprophylaxis in these patients is the avoidance of the small but catastrophic risk of intracranial or spinal bleeding complications associated with pharmacological agents⁹⁷. External

pneumatic compression is effective in these patients^{60,98,99}, although its use may be limited by patient compliance. The ease of application and management of stockings may be an advantage. In a recent randomized controlled trial comparing stockings alone *versus* stockings combined with intermittent pneumatic compression, the DVT rate was 9 per cent for stockings only and 9 per cent for stockings and pneumatic compression, compared with 20 per cent in untreated controls⁶⁰. Stockings alone appeared sufficient. Another multicentre trial showed that LMWH reduced the rate of DVT from 21 per cent with stockings alone, to 14 per cent with stockings and nadroparin (relative risk reduction 34 per cent; $P=0.018$)¹⁰⁰.

Stockings may suffice as the sole agent, especially where pharmacological agents are contraindicated. Graduated compression stockings combined with LMWH or low-dose heparin should provide adequate DVT prophylaxis in neurological and neurosurgical patients.

Medical

In a study of autopsy-proven PE in 2388 hospital deaths, Sandler and Martin¹⁰¹ showed that only 24 per cent of the 239 patients who died from PE had undergone recent surgery. Although the emphasis in the literature is on surgical patients, DVT is also important in medical patients^{102–104}. The efficacy of pharmacological agents, especially heparin, in preventing DVT in patients with heart failure, myocardial infarction and chest infections has been demonstrated by clinical trials^{105,106}. In contrast, clinical evidence of the value of mechanical prophylaxis in medical patients is limited. A recent prospective randomized trial of graduated compression stockings in 80 patients aged 70 years and above with acute myocardial infarction, using the fibrinogen uptake test, showed DVT in

eight control legs compared with none in legs with stockings¹⁰⁷. With the exception of this trial and data on stroke patients, no other studies on the efficacy of graduated compression stockings in general medical patients were found. The available data suggest that there is a place for stockings in DVT prophylaxis in medical patients. This is particularly important when hospitalization is prolonged, in patients aged over 40 years and in the presence of medical conditions known to predispose to DVT.

Recurrent deep vein thrombosis

Previous DVT is a major predisposing factor to DVT recurrence¹⁰⁸. Belcaro *et al.*⁶³ studied the efficacy of graduated compression stockings in preventing recurrent DVT in 224 patients following an episode of DVT. Three years later the recurrence rate was 46 per cent in untreated controls compared with 9, 5 and 2 per cent in patients using stockings, ibuprofen or stockings and ibuprofen respectively.

Comparison with other prophylactic agents

Graduated compression stockings reduce the risk of DVT by 64 per cent in general surgical patients. *Tables 3 and 4* show the results of trials that compare the effect of stockings alone or combined with other prophylactic modalities. Graduated compression stockings combined with low-dose heparin performed better than stockings or heparin alone (*Table 3*). LMWH combined with stockings was better than stockings alone (*Table 4*). Compression stockings and pneumatic compression were equally effective (*Table 5*). Stockings are, however, easier to manage. Either stockings or pneumatic compression may be used during operation and postoperative mechanical prophylaxis can be continued with stockings alone. In hip replacement, a meta-analysis

Table 3 Venous thromboprophylaxis using graduated compression stockings with and without low-dose heparin

Reference	Year	Type of surgery	Diagnostic test	Incidence of deep vein thrombosis		
				LDH	Stockings and LDH	Stockings
Torgren ⁵³	1980	Abdomen	FUT	12 of 98 (12)	4 of 98 (4)	—
Borow and Goldson ¹⁰⁹	1983	Abdomen	FUT	15 of 56 (27)	2 of 63 (3)	—
Wille-Jorgensen <i>et al.</i> ⁵⁵	1985	Abdomen	FUT	11 of 90 (12)	2 of 86 (2)	—
Wille-Jorgensen <i>et al.</i> ⁵⁸	1991	Abdomen	FUT	12 of 81 (15)	2 of 79 (3)	—
Fasting <i>et al.</i> ¹¹⁰	1985	Abdomen	Tc-pl	4 of 45 (9)	3 of 52 (6)	—
Moser <i>et al.</i> ¹¹¹	1976	General	FUT	—	2 of 20 (10)	5 of 20 (25)
Nicolaides <i>et al.</i> ¹¹²	1972	General	FUT	—	1 of 122 (1)	29 of 122 (24)
Borow and Goldson ¹⁰⁹	1983	General	FUT	—	2 of 63 (3)	15 of 106 (14)

Values in parentheses are percentages. LDH, low-dose heparin; FUT, fibrinogen ¹²⁵I uptake test; Tc-pl, Technetium ^{99m}Tc-plasmin scan for deep vein thrombosis

Table 4 Venous thromboprophylaxis using graduated compression stockings with and without low molecular weight heparin

Reference	Year	Type of surgery	Diagnostic test	Incidence of deep vein thrombosis	
				Stockings	Stockings and LMWH
Lassen <i>et al.</i> ¹¹³	1991	Hip arthroplasty	Venography	45 of 97 (46)	30 of 97 (31)
Kalodiki <i>et al.</i> ⁶⁸	1996	Hip arthroplasty	Venography	12 of 32 (38)	8 of 32 (25)
Nurmohamed <i>et al.</i> ¹⁰⁰	1996	Neurosurgery	Venography	51 of 244 (21)	33 of 241 (14)
Levine <i>et al.</i> ¹¹⁴	1996	Knee surgery	Venography	60 of 124 (48)	29 of 96 (30)

Values in parentheses are percentages. LMWH, low molecular weight heparin

Table 5 Venous thromboprophylaxis using graduated compression stockings and pneumatic compression

Reference	Year	Type of surgery	Diagnostic test	Incidence of deep vein thrombosis		
				Stockings	Pneumatic compression	Stockings and pneumatic compression
Bucci <i>et al.</i> ¹¹⁵	1989	Neurosurgery	IPG	1 of 38 (3)	1 of 32 (3)	—
Hansberry <i>et al.</i> ⁹⁵	1991	Urological	Platelet scan*	5 of 25 (20)	3 of 24 (13)	—
Turpie <i>et al.</i> ⁶⁰	1989	Neurosurgery	FUT	7 of 80 (9)	—	7 of 78 (9)
Goldhaber <i>et al.</i> ¹¹⁶	1995	Cardiac	Ultrasonography	38 of 172 (22)	—	33 of 172 (19)

Values in parentheses are percentages. *¹¹¹In-labelled platelet scan. IPG, impedance plethysmography; FUT, fibrinogen ¹²⁵I uptake test

based on 56 trials showed an unadjusted pooled risk of 21 per cent for use of stockings compared with 16 per cent for LMWH and 24 per cent for low-dose heparin⁷⁷. Only LMWH and stockings were shown to reduce the risk of PE in the analysis. Although this agrees with the report by Wilkins *et al.* on the value of stockings in preventing PE, the few trials evaluating this suggestion are limited by small numbers.

Graduated compression stockings alone are effective for the prevention of DVT. Their effect is, however, enhanced by pharmacological agents such as low-dose heparin and LMWH. Stockings alone could provide adequate prophylaxis for patients at low risk of DVT. Patients at moderate or high risk should be managed with stockings combined with LMWH or low-dose heparin^{117,118}.

Thigh-length versus knee-length stockings

The majority of studies on prevention of venous thrombosis have been on thigh- or full-length stockings. Thigh- or full-length stockings are more expensive, more difficult to put on and less tolerated than knee-length stockings^{119–121}. Lawrence and Kakkar⁴³ found no increase in deep venous flow velocity when whole limb compression (18–8 mmHg) was applied compared with below-knee compression (18–14 mmHg). Porteous *et al.*¹²² compared thigh- and knee-

length stockings in a randomized trial using the fibrinogen uptake test and found no significant difference in their ability to prevent DVT. Sparrow *et al.*⁴⁷ observed that the portion of the stocking above the tibial plateau was relatively ineffective in decreasing venous pooling, but appeared to compromise the value of the stocking at lower levels. This prompted concern that the thigh- or full-length stockings may actually be inferior to knee-length stockings. The above-knee segment of a thigh stocking often rolls down, either hanging loosely around the knee or exerting a garter-like tourniquet effect^{121,123}, thus compromising the effect of the stocking and potentially increasing the risk of thrombosis. A recent survey of patient compliance showed a significantly higher number of improperly applied thigh-length than knee-length stockings¹¹⁹. Knee-length stockings should replace thigh-length stockings, being equally effective, cheaper, more likely to fit correctly and better tolerated by patients.

Customized stockings

Customized stockings, tailored to the patient's leg dimensions, are usually made from high-modulus yarn with substantial variations in the pressure profile with changing limb circumference. Johnson *et al.*¹²⁴ compared these with commercially available non-custom low-modulus circular

knit stockings, but found no advantage. Non-customized stockings actually appeared superior in that their low modulus allowed for flexibility in the calf diameter within the target pressure profile. While commercial non-customized stockings may be preferable to tailored stockings, studies suggest that they do not always conform to the pressures specified by their manufacturers^{125,126}.

There may be a case for customized stockings in patients with pronounced calf–ankle disproportion, as in severe lipodermatosclerosis¹²¹.

Indications

Patients at risk of DVT may be classified into low-, moderate- and high-risk groups based on the aggregate of predisposing factors¹³. Graduated compression stockings may be used as sole prophylactic agent in patients at low risk of DVT. Stockings may also be used alone in situations where pharmacological prophylaxis is contraindicated and for long-term prevention of recurrence following 3–6 months of anticoagulation for a single episode of DVT⁶³. Patients at moderate or high risk should be managed with stockings in combination with LMWH or low-dose heparin. Stockings should be applied at least 2 h before operation, and continued during and after surgery until the patient is fully mobile. Stockings are particularly beneficial in the bedridden¹²⁷. The practice of sitting patients out of bed early after operation can be hazardous. Sitting with the knees flexed to 90° or more for a patient with thigh-length stockings causes a marked increase in popliteal–stocking interface pressure. This compromises the effect of the stockings and may jeopardize thromboprophylaxis¹²⁷. The risk of venous thromboembolism subsists after discharge from hospital^{89,128,129}. Stockings may potentially be valuable in reducing this risk. With the increasing trend towards early discharge from hospital, a properly conducted randomized controlled trial of the value of graduated compression stockings following discharge is necessary.

Complications

Although graduated compression stockings are relatively safe, they are not without risk. The main concern is impairment of subcutaneous tissue oxygenation, especially in patients with existing peripheral vascular compromise. A stocking pressure of 10 mmHg produces a 10 per cent reduction in cutaneous blood flow, 30 mmHg a 25 per cent drop and 60 mmHg an 84 per cent drop¹³⁰. An estimate of the hazards of compression treatment of the leg by 154 Scottish surgeons showed that at some stage up to one-third had recognized a patient with damage to the leg as a result of

compression therapy¹³¹. Compression stockings for DVT prophylaxis were associated with 38 of the 147 cases reported. Patients with peripheral arterial disease and diabetics with neuropathy are particularly at risk, and cases resulting in necrosis and amputation have been reported^{132,133}. Arterial occlusion, thrombosis and gangrene over the anterior aspect of the ankle secondary to the tourniquet effect of multiple layers of rolled down stocking have also been reported^{123,134}. It is therefore essential that significant arterial compromise and peripheral neuropathy are excluded before applying stockings. Manufacturers advise against the use of stockings in patients with an ankle:brachial pressure index less than 0.7. Stockings should also be sized and fitted correctly. Leg measurements should be reviewed regularly especially in swollen or swelling legs, as changes in leg girth significantly change the amount of pressure exerted rendering the stocking ineffective or threatening tissue oxygenation. An increase in leg circumference of 5 cm can double the amount of pressure being applied by a stocking¹³⁵.

Conclusions

Graduated compression stockings provide an effective, safe, cheap and convenient means of preventing DVT. Stockings increase the linear velocity of venous outflow, prevent stasis and venous distension, and enhance emptying of valvular cusps. Properly used, stockings can reduce the risk of DVT in hospitalized patients by 55–70 per cent. Evidence on the value of graduated compression stockings in preventing fatal PE is limited and inconclusive. In combination with LMWH or low-dose heparin, stockings provide adequate prophylaxis for the majority of patients at risk. Knee-length stockings are preferable, being effective and potentially superior to thigh- or full-length stockings. Knee-length stockings are cheaper and better tolerated, and should replace thigh- and full-length stockings for DVT prophylaxis. Continued use of stockings to prevent late venous thrombosis following discharge from hospital may be beneficial in patients with poor mobility. The role of stockings in preventing DVT recurrence is noteworthy. Attention to the vascular and neurological status of the leg, proper sizing and regular review of the legs should limit the rare occurrence of ischaemic complications with stockings.

Acknowledgements

The authors are grateful to the Cochrane group, whose database was searched for studies relevant to this review.

References

- 1 Cohen AT, Edmondson RA, Phillips MJ, Kakkar VV. The changing pattern of thromboembolic disease. *Haemostasis* 1996; **26**: 65–71.
- 2 Collins R, Scrimgeour A, Yusuf S, Peto R. Reduction in fatal pulmonary embolism and venous thrombosis by perioperative administration of subcutaneous heparin. Overview of results of randomized trials in general, orthopedic, and urologic surgery. *N Engl J Med* 1988; **318**: 1162–73.
- 3 Dismuke SE, Wagner EH. Pulmonary embolism as a cause of death. The changing mortality in hospitalised patients. *JAMA* 1986; **255**: 2039–42.
- 4 Lilienfeld DE, Chan E, Ehland J, Godbold JH, Landrigan PJ, Marsh G. Mortality from pulmonary embolism in the United States: 1962 to 1984. *Chest* 1990; **98**: 1067–72.
- 5 National Institute of Health Consensus Conference. Prevention of venous thrombosis and pulmonary embolism. *JAMA* 1986; **256**: 744–9.
- 6 Colditz GA, Tuden RL, Oster G. Rates of venous thrombosis after general surgery: combined results of randomised clinical trials. *Lancet* 1986; **ii**: 143–6.
- 7 Hull R, Raskob GE, Hirsh J. Prophylaxis of venous thromboembolism: an overview. *Chest* 1986; **89**(Suppl 5): 374–83s.
- 8 Clagett GP, Reisch JS. Prevention of venous thromboembolism in general surgical patients. Results of meta-analysis. *Ann Surg* 1988; **208**: 227–40.
- 9 Stamatakis JD, Sagar S, Maffei FH, Lawrence D, Kakkar VV. Femoral vein thrombosis and total hip arthroplasty. *Br J Surg* 1976; **63**: 668–9 (Abstract).
- 10 Stulberg BN, Insall JN, Williams GW, Ghelman B. Deep-vein thrombosis following total knee replacement. An analysis of six hundred and thirty-eight arthroplasties. *J Bone Joint Surg Am* 1984; **66**: 194–201.
- 11 Jeffery PC, Nicolaides AN. Graduated compression stockings in the prevention of postoperative deep vein thrombosis. *Br J Surg* 1990; **77**: 380–3.
- 12 Parker-Williams J, Vickers R. Major orthopaedic surgery on the leg and thromboembolism. *BMJ* 1991; **303**: 531–2.
- 13 Thromboembolic Risk Factors (THRIFT) Consensus Group. Risk and prophylaxis for venous thromboembolism in hospital patients. *BMJ* 1992; **305**: 567–74.
- 14 Caprini JA, Chucker JL, Zuckerman L, Vagher JP, Frank CA, Cullen JE. Thrombosis prophylaxis using external compression. *Surg Gynecol Obstet* 1983; **156**: 599–604.
- 15 Bergqvist D. Prophylaxis against postoperative deep vein thrombosis: a survey of surveys. *Thromb Haemost* 1990; **2**: 69–73.
- 16 Fletcher JP, Koutris J, Ockelford PA. Deep vein thrombosis prophylaxis: a survey of current practice in Australia and New Zealand. *Aust NZ J Surg* 1992; **62**: 601–5.
- 17 Anderson FA Jr, Wheeler HB, Goldberg RJ, Hosmer DW, Forcier A, Patwardhan NA. Physician practices in the prevention of venous thromboembolism. *Ann Intern Med* 1991; **115**: 591–5.
- 18 George BD, Cook TA, Franklin IJ, Nethercliff J, Galland RB. Protocol violation in deep vein thrombosis prophylaxis. *Ann R Coll Surg Engl* 1998; **80**: 55–7.
- 19 Hui ACW, Heras-Palou C, Dunn I, Triffitt PD, Crozier A, Imeson J *et al*. Graded compression stockings for the prevention of deep vein thrombosis after hip and knee replacement. *J Bone Joint Surg Br* 1996; **78-B**: 550–4.
- 20 Ohlund C, Fransson SG, Starck SA. Calf compression for prevention of thromboembolism following hip surgery. *Acta Orthop Scand* 1983; **54**: 896–9.
- 21 Nilsen DWT, Naess-Andresen KF, Kierulf P, Heldaas J, Storen G, Godal HC. Graded pressure stockings in prevention of deep vein thrombosis following total hip replacement. *Acta Chirurgica Scandinavica* 1984; **150**: 531–4.
- 22 Stanton JR, Freis ED, Wilkins RW. The acceleration of linear flow in the deep veins of the lower extremity of man by local compression. *J Clin Invest* 1949; **28**: 553–8.
- 23 Meyerowitz BR, Nelson R. Measurement of the velocity of blood in lower limb veins with and without compression. *Surgery* 1964; **56**: 481–6.
- 24 Wright HP, Osborn SB. Effect of posture on venous velocity, measured with Na²⁴Cl. *Br Heart J* 1952; **14**: 325–30.
- 25 Spiro M, Roberts VC, Richards JB. Effects of externally applied pressure on femoral vein blood flow. *BMJ* 1970; **i**: 719–23.
- 26 Ido K, Suzuki T, Taniguchi Y, Kawamoto C, Isoda N, Nagamine N *et al*. Femoral vein stasis during laparoscopic cholecystectomy: effects of graded elastic compression leg bandages in preventing thrombus formation. *Gastrointest Endosc* 1995; **42**: 151–5.
- 27 Sigel B, Edelstein AL, Felix WR Jr, Memhardt CR. Compression of the deep venous system of the leg during inactive recumbency. *Arch Surg* 1973; **106**: 38–43.
- 28 Litter J. Thromboembolism: its prophylaxis and medical treatment. *Med Clin North Am* 1952; **36**: 1309–21.
- 29 Lewis CE Jr, Antoine J, Mueller C, Talbot WA, Swaroop R, Edwards WS. Elastic compression in the prevention of venous stasis. A critical reevaluation. *Am J Surg* 1976; **132**: 739–43.
- 30 McLachlin AD, McLachlin JA, Jory TA, Rawling EG. Venous stasis in the lower extremities. *Ann Surg* 1960; **152**: 678–85.
- 31 Sarin S, Scurr JH, Coleridge Smith PD. Mechanism of action of external compression on venous function. *Br J Surg* 1992; **79**: 499–502.
- 32 Comerota AJ, Stewart GJ, Alburger PD, Smalley K, White JV. Operative venodilation: a previously unsuspected factor in the cause of postoperative deep vein thrombosis. *Surgery* 1990; **106**: 301–9.
- 33 Coleridge Smith PD, Hasty JH, Scurr JH. Venous stasis and lumen changes during surgery. *Br J Surg* 1990; **77**: 1055–9.
- 34 Stewart GJ, Schaub RG, Niewiarowski S. Products of tissue injury. Their induction of venous endothelial damage and

- blood cell adhesion in the dog. *Arch Pathol Lab Med* 1980; **104**: 409–13.
- 35 Coleridge Smith PD, Hasty JH, Scurr JH. Deep vein thrombosis: effect of graduated compression stockings on distension of the deep veins of the calf. *Br J Surg* 1991; **78**: 724–6.
- 36 Arcelus JI, Caprini JA, Hoffman KN, Traverso CI, Hoppensteadt D, Fareed J *et al*. Modifications of plasma levels of tissue factor pathway inhibitor and endothelin-1 induced by reverse Trendelenburg position: influence of elastic compression – preliminary results. *J Vasc Surg* 1995; **22**: 568–72.
- 37 Hansen JB, Olsen JO, Osterud B. Physical exercise enhances the plasma levels of extrinsic pathway inhibitor (EPI). *Thromb Haemost* 1990; **64**: 124–6.
- 38 Graboski EF, Zuckerman DB, Nemerson Y. The functional expression of tissue factor by fibroblasts and endothelial cells under flow conditions. *Blood* 1993; **81**: 3265–70.
- 39 Holst J, Lindblad B, Wedeberg E, Bergqvist D, Nordfang O, Ostergaard PB *et al*. Tissue factor pathway inhibitor (TFPI) and its response to heparin in patients with spontaneous deep vein thrombosis. *Thromb Res* 1993; **72**: 467–70.
- 40 Kijowski R, Hoppensteadt DA, Walenga J, Borris L, Lassen MR, Fareed J. Role of tissue factor pathway inhibitor in post surgical deep vein thrombosis (DVT) prophylaxis in patients treated with low molecular weight heparin. *Thromb Res* 1994; **74**: 53–64.
- 41 Walenga JM, Koza MJ, Lewis BE *et al*. Relative heparin induced thrombocytopenic potential of low molecular weight heparins and new antithrombotic agents. *Thromb Haemost* 1996; **2**(Suppl 1): S21.
- 42 Sigel B, Edelstein AL, Savitch L, Hasty JH, Felix WR Jr. Type of compression for reducing venous stasis. A study of lower extremities during inactive recumbency. *Arch Surg* 1975; **110**: 171–5.
- 43 Lawrence D, Kakkar VV. Graduated, static, external compression of the lower limb: a physiological assessment. *Br J Surg* 1980; **67**: 119–21.
- 44 Holford CP. Graded compression for preventing deep vein thrombosis. *BMJ* 1976; **ii**: 969–70.
- 45 Allan A, Williams JT, Bolton JP, Le Quesne LP. The use of graduated compression stockings in the prevention of postoperative deep vein thrombosis. *Br J Surg* 1983; **70**: 172–4.
- 46 Gaylarde PM, Sarkany I, Dodd HJ. The effect of compression on venous stasis. *Br J Dermatol* 1993; **128**: 255–8.
- 47 Sparrow RA, Hardy JG, Fentem PH. Effect of ‘antiembolism’ compression hosiery on leg blood. *Br J Surg* 1995; **82**: 53–9.
- 48 Wilkins RW, Mixer G Jr, Stanton JR, Litter J. Elastic stockings in the prevention of pulmonary embolism: a preliminary report. *N Engl J Med* 1952; **246**: 360–4.
- 49 Makin GS. A clinical trial of ‘Tubigrip’ to prevent deep venous thrombosis. *Br J Surg* 1969; **56**: 373–5.
- 50 Rosengarten DS, Laird J, Jeyasingh K, Martin P. The failure of compression stockings (Tubigrip) to prevent deep venous thrombosis after operation. *Br J Surg* 1970; **57**: 296–9.
- 51 Browse NL, Jackson BT, Mayo ME, Negus D. The value of mechanical methods of preventing postoperative calf vein thrombosis. *Br J Surg* 1974; **61**: 219–23.
- 52 Scurr JH, Ibrahim SZ, Faber RG, Le Quesne LP. The efficacy of graduated compression stockings in the prevention of deep vein thrombosis. *Br J Surg* 1977; **64**: 371–3.
- 53 Tornngren S. Low dose heparin and compression stockings in the prevention of postoperative deep venous thrombosis. *Br J Surg* 1980; **67**: 482–4.
- 54 Bergqvist D, Lindblad B. The thromboprophylactic effect of graded elastic compression stockings in combination with dextran 70. *Arch Surg* 1984; **119**: 1329–31.
- 55 Wille-Jorgensen P, Thorup J, Fischer A, Holst-Christensen J, Flamsholt R. Heparin with and without graded compression stockings in the prevention of thromboembolic complications of major abdominal surgery: a randomized trial. *Br J Surg* 1985; **72**: 579–81.
- 56 Mellbring G, Palmer K. Prophylaxis of deep vein thrombosis after major abdominal surgery. Comparison between dihydroergotamine–heparin and intermittent calf compression and evaluation of added graduated static compression. *Acta Chirurgica Scandinavica* 1986; **152**: 597–600.
- 57 Scurr JH, Coleridge Smith PD, Hasty JH. Regimen for improved effectiveness of intermittent pneumatic compression in deep venous thrombosis prophylaxis. *Surgery* 1987; **102**: 816–20.
- 58 Wille-Jorgensen Hauch O, Dimo B, Christensen SW, Jensen R, Hansen B. Prophylaxis of deep venous thrombosis after acute abdominal operation. *Surg Gynecol Obstet* 1991; **172**: 44–8.
- 59 Turner GM, Cole SE, Brooks JH. The efficacy of graduated compression stockings in the prevention of deep venous thrombosis after major gynaecological surgery. *Br J Obstet Gynaecol* 1984; **91**: 588–91.
- 60 Turpie AG, Hirsh J, Gent M, Julian D, Johnson J. Prevention of deep vein thrombosis in potential neurosurgical patients. A randomized trial comparing graduated compression stockings alone or graduated compression stockings plus intermittent pneumatic compression with control. *Arch Intern Med* 1989; **149**: 679–81.
- 61 Wells PS, Lensing AWA, Hirsh J. Graduated compression stockings in the prevention of postoperative venous thromboembolism. A meta-analysis. *Arch Intern Med* 1994; **154**: 67–72.
- 62 Markel A, Manzo RA, Bergelin RO, Strandness DE Jr. Valvular reflux after deep vein thrombosis: incidence and time of occurrence. *J Vasc Surg* 1992; **15**: 377–84.
- 63 Belcaro G, Laurora G, Cesarone MR, De Sanctis MT. Prophylaxis of recurrent deep venous thrombosis. A randomized, prospective study using indobufen and graduated elastic compression stockings. *Angiology* 1993; **44**: 695–9.
- 64 Brandjes DPM, Buller HR, Heijboer H, Huisman MV, de Rijk M, Jagt H *et al*. Randomised trial of effect of compression

- stockings in patients with symptomatic proximal-vein thrombosis. *Lancet* 1997; **349**: 759–62.
- 65 Morris GK. Prevention of venous thromboembolism. A survey of methods used by orthopaedic and general surgeons. *Lancet* 1980; **ii**: 572–4.
- 66 Ishak MA, Morley KD. Deep venous thrombosis after total hip arthroplasty: a prospective controlled study to determine the prophylactic effect of graded pressure stockings. *Br J Surg* 1981; **68**: 429–32.
- 67 Fredin H, Bergqvist D, Cederholm C, Lindblad B, Nyman U. Thromboprophylaxis in hip arthroplasty. Dextran with graded compression or preoperative dextran compared in 150 patients. *Acta Orthop Scand* 1989; **60**: 678–81.
- 68 Kalodiki EP, Hoppensteadt DA, Nicolaidis AN, Fareed J, Gill K, Regan F *et al.* Deep venous thrombosis prophylaxis with low molecular weight and elastic compression in patients having total hip replacement. A randomised controlled trial. *Int Angiol* 1996; **15**: 162–8.
- 69 Barnes RW, Brand RA, Clarke W, Hartley N, Hoak JC. Efficacy of graded-compression antiembolism stockings in patients undergoing total hip arthroplasty. *Clin Orthop* 1978; **132**: 61–7.
- 70 Bergqvist E, Bergqvist D, Bronge A, Dahlgren S, Lindquist B. An evaluation of early thrombosis prophylaxis following fracture of the femoral neck. A comparison between Dextran and Dicumarol. *Acta Chirurgica Scandinavica* 1972; **138**: 689.
- 71 Beisaw NE, Camerota AJ, Groth HE, Merli GJ, Weitz HH, Zimmerman RC *et al.* Dihydroergotamine/heparin in the prevention of deep vein thrombosis after total hip replacement. A controlled, prospective, randomized multicenter trial. *J Bone Joint Surg Am* 1988; **70**: 2–10.
- 72 Salzman EW, Harris WH. Prevention of venous thromboembolism in orthopaedic patients. *J Bone Joint Surg Am* 1976; **58**: 903–12.
- 73 Everts CM, Feil EJ. Prevention of thromboembolic disease after elective surgery of the hip. *J Bone Joint Surg Am* 1971; **53**: 1271–80.
- 74 Laverick MD, Croal SA, Mollan RAB. Orthopaedic surgeons and thromboprophylaxis. *BMJ* 1991; **303**: 549–50.
- 75 Nicolaidis AN, Miles C, Hoare M, Jury P, Helms E, Venniker R. Interim sequential pneumatic compression of the legs and thromboembolism-deterrent stockings in the prevention of postoperative deep venous thrombosis. *Surgery* 1983; **94**: 21–5.
- 76 Cotton LT, Roberts VC. The prevention of deep vein thrombosis, with particular reference to mechanical methods of prevention. *Surgery* 1977; **81**: 228–35.
- 77 Imperiale TF, Speroff T. A meta-analysis of methods to prevent venous thromboembolism following total hip replacement. *JAMA* 1994; **271**: 1780–5.
- 78 Hirsh J. Low-molecular-weight heparin for the treatment of venous thromboembolism. *Am Heart J* 1998; **135**: S336–42.
- 79 Hull R, Raskob G, Pineo G, Rosenbloom D, Evans W, Mallory T *et al.* A comparison of subcutaneous low-molecular-weight heparin with warfarin sodium for prophylaxis against deep-vein thrombosis after hip and knee implantation. *N Engl J Med* 1993; **329**: 1370–6.
- 80 Hoek JA, Nurmohamed MT, Hamelynck KJ, Marti RK, Knipscheer HC, ten Cate H *et al.* Prevention of deep vein thrombosis following total hip replacement by low molecular weight heparinoid. *Thromb Haemost* 1992; **67**: 28–32.
- 81 Spiro TE, Colwell CW, Bona RD, Trowbridge AA, Wise GR, Furman KW *et al.* A clinical trial comparing the efficacy and safety of enoxaparin, a low molecular weight heparin and unfractionated heparin for the prevention of deep venous thrombosis after elective knee replacement surgery. *Blood* 1993; **82**(Suppl): 410a (Abstract).
- 82 Leclerc JR, Geerts WH, Desjardins L, L'Espérance M, Cruickshank M, Demers C *et al.* Prevention of venous thromboembolism after knee arthroplasty: a randomized double-blind trial, comparing a low molecular weight heparin fragment (enoxaparin) to warfarin. *Blood* 1994; **82**(Suppl): 246a (Abstract).
- 83 Stamatakis JD, Kakkar VV, Sagar S, Lawrence D, Nairn D, Bentley PG. Femoral vein thrombosis and total hip replacement. *BMJ* 1977; **ii**: 223–5.
- 84 Merli GJ. Deep vein thrombosis and pulmonary embolism prophylaxis in orthopedic surgery: update. *Med Clin North Am* 1993; **77**: 397–411.
- 85 Abdel-Salam A, Eyres KS. Effects of tourniquet during total knee arthroplasty. A prospective randomised study. *J Bone Joint Surg Br* 1995; **77-B**: 250–3.
- 86 Wakankar HM, Nicholl JE, Koka R, D'Arcy JC. The tourniquet in total knee arthroplasty. A prospective, randomised study. *J Bone Joint Surg Br* 1999; **81-B**: 30–3.
- 87 Parmet JL, Berman AT, Horrow JC, Harding S, Rosenberg H. Thromboembolism coincident with tourniquet deflation during total knee arthroplasty. *Lancet* 1993; **341**: 1057–8.
- 88 Parmet JL, Horrow JC, Berman AT, Miller F, Pharo G, Collins L. The incidence of large venous emboli during total knee arthroplasty without pneumatic tourniquet use. *Anesth Analg* 1998; **87**: 439–44.
- 89 Planes A, Vochelle N, Darmon JY, Fagola M, Bellaud M, Huet Y. Risk of deep-venous thrombosis after hospital discharge in patients having undergone total hip replacement: double-blind randomised comparison of enoxaparin *versus* placebo. *Lancet* 1996; **348**: 224–8.
- 90 Trowbridge A, Boese CK, Woodruff B, Brindley HH Sr, Lowry WE, Spiro TE. Incidence of post hospitalization proximal deep vein thrombosis after total hip arthroplasty. A pilot study. *Clin Orthop* 1994; **299**: 203–8.
- 91 Collins REC, Klein LA, Skillman JJ, Salzman EW. Thromboembolic problems in urologic surgery. *Urol Clin North Am* 1976; **3**: 393–401.
- 92 Walsh JJ, Bonnar J, Wright FW. A study of pulmonary embolism after deep vein thrombosis after major gynaecological surgery using labelled fibrinogen-plebography and lung scanning. *J Obstet Gynaecol Br Commonwealth* 1974; **81**: 311–16.
- 93 Hobel CJ, Castro L, Rosen D, Greenspoon JS, Nessim S. The effect of thigh-length support stockings on the hemodynamic

- response to ambulation in pregnancy. *Am J Obstet Gynecol* 1996; **174**: 1734–41.
- 94 Nilsson L, Austrell CH, Norgren L. Venous function during late pregnancy. The effect of elastic compression hosiery. *Vasa* 1992; **21**: 203–5.
- 95 Hansberry KL, Thompson IM Jr, Bauman J, Deppe S, Rodriguez FR. A prospective comparison of thromboembolic stockings, external sequential pneumatic compression stockings and heparin sodium/dihydroergotamine mesylate for the prevention of thromboembolic complications in urological surgery. *J Urol* 1991; **145**: 1205–8.
- 96 Hamilton MG, Hull RD, Pineo GF. Venous thromboembolism in neurosurgery and neurology patients: a review. *Neurosurgery* 1994; **34**: 280–96.
- 97 Cerrato D, Ariano C, Fiacchino F. Deep vein thrombosis and low-dose heparin prophylaxis in neurosurgical patients. *J Neurosurg* 1978; **48**: 378–81.
- 98 Turpie AGG, Gallus A, Beattie WS, Hirsh J. Prevention of venous thrombosis in patients with intracranial disease by intermittent pneumatic compression of the calf. *Neurology* 1977; **27**: 435–8.
- 99 Skillman JJ, Collins REC, Coe NP, Goldstein BS, Shapiro RM, Zervas NT *et al.* Prevention of deep vein thrombosis in neurosurgical patients: a controlled randomised trial of external pneumatic compression boots. *Surgery* 1978; **83**: 354–8.
- 100 Nurmohamed MT, van Riel AM, Henkens CMA, Koopman MM, Que GT, d'Azemar P *et al.* Low molecular weight heparin and compression stockings in the prevention of venous thromboembolism in neurosurgery. *Thromb Haemost* 1996; **75**: 233–8.
- 101 Sandler DA, Martin JF. Autopsy proven pulmonary embolism in hospital patients: are we detecting enough deep vein thrombosis? *J R Soc Med* 1989; **82**: 203–5.
- 102 Nicolaides AN, Irving D. Clinical factors and the risk of deep venous thrombosis. In: Nicolaides AN, eds. *Thromboembolism: Aetiology, Advances in Prevention and Management*. Lancaster: MTP, 1975: 193–204.
- 103 Nicolaides AN, Kakkar VV, Renney JTG, Kidner PH, Hutchison DC, Clarke MB. Myocardial infarction and deep-vein thrombosis. *BMJ* 1971; **i**: 432–4.
- 104 Coon WW. Epidemiology of venous thromboembolism. *Ann Surg* 1977; **186**: 149–64.
- 105 Harenberg J, Kallenbach B, Martin U, Dempfle CE, Zimmermann R, Kubler W *et al.* Randomized controlled trial of heparin and low molecular weight heparin for prevention of deep-vein thrombosis in medical patients. *Thromb Res* 1990; **59**: 639–50.
- 106 Belch JJ, Lowe GDO, Ward AG, Forbes CD, Prentice CR. Prevention of deep vein thrombosis in medical patients by low-dose heparin. *Scott Med J* 1981; **26**: 115–17.
- 107 Keiekegaard A, Norgren L. Graduated compression stockings in the prevention of deep vein thrombosis in patients with acute myocardial infarction. *Eur Heart J* 1993; **14**: 1365–8.
- 108 Meissner MH, Caps MT, Bergelin RO, Manzo RA, Strandness DE Jr. Propagation, rethrombosis and new thrombus formation after acute deep vein thrombosis. *J Vasc Surg* 1995; **22**: 558–67.
- 109 Borow M, Goldson H. Prevention of postoperative deep venous thrombosis and pulmonary emboli with combined modalities. *Am Surg* 1983; **49**: 599–605.
- 110 Fasting H, Anderson K, Nielsen HK, Husted SE, Koopmann HD, Simonsen O *et al.* Prevention of postoperative deep venous thrombosis. Low-dose heparin *versus* graded pressure stockings. *Acta Chirurgica Scandinavica* 1985; **151**: 245–8.
- 111 Moser G, Froidevaux A, Nicolaides AN, Dupont PA, Desai S, Lewis JD *et al.* Small doses of subcutaneous sodium heparin in preventing deep venous thrombosis after major surgery. *Schweiz Rundsch Med Prax* 1976; **65**: 1015–20.
- 112 Nicolaides AN, Dupont PA, Desai S, Lewis JD, Douglas JN, Dodsworth H *et al.* Small doses of subcutaneous heparin in preventing deep venous thrombosis after major surgery. *Lancet* 1972; **ii**: 890–3.
- 113 Lassen MR, Borris LC, Christiansen HM, Boll KL, Eiskjær SP, Nielsen BW *et al.* Prevention of thromboembolism in 190 hip arthroplasties. Comparison of LMW heparin and placebo. *Acta Orthop Scand* 1991; **62**: 33–8.
- 114 Levine MN, Gent M, Hirsh J, Weitz J, Turpie AG, Powers P *et al.* Ardeparin (low-molecular-weight heparin) *vs* graduated compression stockings for the prevention of venous thromboembolism. A randomized trial in patients undergoing knee surgery. *Arch Intern Med* 1996; **156**: 851–6.
- 115 Bucci MN, Papadopoulos SM, Chen JC, Campbell JA, Hoff JT. Mechanical prophylaxis of venous thrombosis in patients undergoing craniotomy: a randomized trial. *Surg Neurol* 1989; **32**: 285–8.
- 116 Goldhaber SZ, Hirsch DR, MacDougall RC, Polak JF, Creager MA, Cohn LH. Prevention of venous thrombosis after coronary artery bypass surgery (a randomised trial comparing two mechanical prophylaxis strategies). *Am J Cardiol* 1995; **76**: 993–6.
- 117 Wille-Jørgensen P. *Prophylaxis of Postoperative Thromboembolism*. Copenhagen: Laegeforeningens Forlag, 1991.
- 118 Prevention of venous thromboembolism. International Consensus Statement. *Int Angiol* 1997; **16**: 3–38.
- 119 Williams AM, Davies PR, Sweetnam DIS, Harper G, Pusey R, Lightowler CDR. Knee-length versus thigh-length graduated stockings in the prevention of deep vein thrombosis. *Br J Surg* 1996; **83**: 1553.
- 120 Williams JT, Palfery SM. Cost effectiveness of below knee against above knee compression stockings in the prevention of deep vein thrombosis. *Phlebologie* 1988; **41**: 809–11.
- 121 Whitley AJ. Elastic stockings. *BMJ* 1988; **296**: 413–14.
- 122 Porteous LMJ, Nicholson EA, Morris LT, James R, Negus D. Thigh length versus knee length stockings in the prevention of deep vein thrombosis. *Br J Surg* 1989; **76**: 296–7.
- 123 Heath DI, Kent SJS, Johns DL, Young TW. Arterial

- thrombosis associated with graduated pressure antiembolic stockings. *BMJ* 1987; **295**: 580.
- 124 Johnson G Jr, Kupper C, Farrar DJ, Swallow RT. Graded compression stockings; custom *vs* noncustom. *Arch Surg* 1982; **117**: 69–72.
- 125 Williamson M, Thomas S, Edwards A, Johnson R, Riggs J, Lewis MH. Graduated compression stockings in the prevention of postoperative deep vein thrombosis: a comparative study of pressure profiles and patient compliance. *Phlebology* 1990; **5**: 135–9.
- 126 Thomas S. Anti-embolism stockings. *Pharmaceutical Journal* 1989; **243**: 633.
- 127 Wildin CJ, Hui ACW, Esler CNA, Gregg PJ. *In vivo* pressure profiles of thigh-length graduated compression stockings. *Br J Surg* 1998; **85**: 1228–31.
- 128 Scurr JH, Coleridge Smith PD, Hasty JH. Deep vein thrombosis; a continuing problem. *BMJ* 1988; **297**: 28.
- 129 Huber O, Bounameaux H, Borst F, Rohner A. Postoperative pulmonary embolism after hospital discharge. An underestimated risk. *Arch Surg* 1992; **127**: 310–13.
- 130 Halperin MH, Friedland CK, Wilkins RW. The effect of local compression upon blood flow in the extremities of man. *Am Heart J* 1948; **35**: 221–37.
- 131 Callam MJ, Ruckley CV, Dale JJ, Harper DR. Hazards of compression treatment of the leg: an estimate from Scottish surgeons. *BMJ* 1987; **295**: 1382.
- 132 Merrett ND, Hanel KC. Ischaemic complications of graduated compression stockings in the treatment of deep venous thrombosis. *Postgrad Med J* 1993; **69**: 232–4.
- 133 Kay TWH, Martin FI. Heel ulcers in patients with long-standing diabetes who wear antiembolism stockings. *Med J Aust* 1986; **145**: 290–1.
- 134 Tillman RM, Marsh DR, Leyland H. Minerva. *BMJ* 1991; **303**: 1560.
- 135 Horner J, Lowth LC, Nicolaides AN. A pressure profile for elastic stockings. *BMJ* 1980; **280**: 818–20.